

FIG. 1




FIG. 2




FIG. 3




Figure 1 consists of 12 bar charts, labeled (a) through (l), each representing a different fish species. The y-axis for all charts is 'Percentage of total catch' ranging from 0 to 100. The x-axis for all charts is 'Year' with values 1990, 1991, 1992, 1993, and 1994. The species and their approximate percentage values are as follows:

- (a) Atlantic croaker: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)
- (b) Striped bass: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)
- (c) Weakfish: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)
- (d) Spot: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)
- (e) Blue crab: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)
- (f) Rockfish: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)
- (g) Atlantic silverside: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)
- (h) Atlantic herring: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)
- (i) Atlantic menhaden: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)
- (j) Atlantic bluefish: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)
- (k) Atlantic tomcod: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)
- (l) Atlantic sand lance: 1990 (~10%), 1991 (~10%), 1992 (~10%), 1993 (~10%), 1994 (~10%)




FIG. 5

